



# Gregor Mendel



## The 'discovery' of modern genetics

Gregor Johann Mendel (1822–1884) was an Austrian monk who closely studied the patterns of inheritance in pea plants.

Click "**start**" to find out more about Mendel, and how his discoveries lead to our understanding of modern genetics.



start



# Mendel's experiments

Over seven years, Mendel experimented on more than 28,000 pea plants! Why were his experiments so successful?

- Pea plants grow quickly.
- Pea plants are available in pure-bred (**homozygous**) strains.
- Many pea plant characteristics show discontinuous variation; they are either one form or another, with no intermediates. This means that their phenotypes are easily distinguishable.




Jupiterimages Corporation

## Mendel's early experiments

### Experiment 1: Cross-pollination

First, Mendel cross-pollinated a homozygous (pure-bred) smooth pea plant and a homozygous wrinkly pea plant.

What type of peas were produced by this cross?



The type of experiment that Mendel carried out, investigating just a single characteristic, is called a **monohybrid cross**.

There are two alleles controlling pea shape. This means there are **three** possible genotypes that the F<sub>2</sub> generation of plants could inherit, leading to **two** possible phenotypes.

	<b>Genotype</b>	<b>Phenotype</b>
homozygous dominant	SS	smooth
homozygous recessive	ww	wrinkly
heterozygous	Sw	smooth

The likelihood of a trait being produced during a monohybrid cross can be mapped out using a **Punnett Square**.



## What do Punnett Squares show?

Punnett Squares show each of the possible outcomes of a monohybrid cross.

Click "**play**" to find out how Punnett Squares explain the 3:1 ratio of smooth to wrinkly peas.



smooth

x



wrinkly



After his research, Mendel proposed two laws of inheritance.

## **Mendel's first law: the law of segregation**

- Alternate versions of genes (alleles) cause variation in inherited characteristics.
- An organism inherits two alleles for each characteristic – one from each parent.
- Dominant alleles will always mask recessive alleles.
- The two alleles for each characteristic separate during gamete production.

## **Mendel's second law: law of independent assortment**

- Genes for different characteristics are sorted independently during gamete production.

